A STUDY ON SOME MORPHOMETRIC TRAITS OF RED SOKOTO AND SAHEL GOATS IN MAIGATARI LOCAL GOVERNMENT AREA OF JIGAWA STATE

Adamu, H., Ma'aruf, B. S., Shuaibu, A. & Umar, H. A
Department of Animal Science, Faculty of Agriculture, Federal University of Kashere, Gombe P.M.B 0182 Gombe State
Corresponding author email: *habibuadamu2016@gmail.com*

ABSTRACT

The study was conducted in Maigatari Local Government Area of Jigawa State. A total of 250 goats (187 Red Sokoto and 63 Sahel) of mixed sex and age were used to obtain the following parameters; body weight (BW), body condition score (BCS), body length (BL), heart girth (HG), wither height (WH), udder length (UL), udder circumference (UC), testicular length (TL) and testicular circumference (TC). The animals were sampled from Maigatari main livestock market. Data generated were subjected to analysis of variance, Pearson's correlation analysis and stepwise linear regression. Red Sokoto had mean BW, BCS, BL, HG, WH, UL, UC, TL and TC of 20.58 kg, 3.20, 22.45 cm, 25.28 cm, 22.93 cm, 3.48 cm, 5.99 cm, 4.97 cm and 7.88 cm, respectively. The corresponding values for Sahel were 22.91 kg, 3.00, 23.58 cm, 25.07 cm, 24.55 cm, 4.14 cm, 7.06 cm, 4.75 cm and 7.52 cm. Sahel goats had significantly higher BL (P<0.01) and WH (P<0.001). Female goats were heavier than male and had longer body, wider chest and higher height at wither. Except TC, goats of 36 - 48 months had the highest for all the significantly affected traits (BW, BCS, BL, HG and WH; P<0.001). The correlation coefficients observed among the morphometric traits were in general high, positive and significant, while the accuracy of the predictive equation when HG and in combination with BL were used to predict live weight was also high. This study indicated that breed had no effect on most body characters and that live weight can best be predicted using HG.

Key words: Morphometrics traits, breed, sex, age, goat.

INTRODUCTION

Small ruminants are important genetic resources and play apredominant role in the sustenance of the livelihoods of impoverished families especially inthe rural areas of tropical countries. Caprine species specifically goats play animportant role in the social life of many Africans and supply human population withmeat, milk, skin, manure and other products. The Red Sokoto goat (RSG) or Maradi is the most predominant breed and accountsfor about 70% of Nigeria's total goat population (Ademosun, 1994). It is commonly found with the agro pastoralist mainly within thenorthern sub-humid and semi-arid zones of the country (Akpa *et al.,* 2001). The Sahel goat is an extant meat and milk type goat in Nigeria (Otoikhian and Orheruata, 2010). It is large in size, predominantly white colour, pied with black or white and brown spots around the ear, nose and udder. This breed is found alongthe northern border of Nigeria, particularly Borno State, where it is often known as"Balami, although this name has not been adopted as it would lead to confusion with thebetter known sheep breed Balami. Morphometric characteristics are important in breed identification and classification. Gizaw *et al.* (2007) stated that morphostructural description of traits remained essential component of characterization that can be used to physically identify, describe and recognize a breed. It also helps in the classification of livestock breeds. The knowledge of morphometric traits aids in appropriate breeding design, feeding and health management (Thiruvenkadan, 2005).

MATERIALS AND METHODS

Study Location

The study was conducted at Maigatari Local Area of Jigawa State. Maigatari is located between longitude 48.26^0 12' N and latitude 25.50^0 9'E and is 352m above sea level (Encarta Encyclopedia, 2009 PC version). It has an average annual rainfall of 418 mm per annum, which starts from late April or early May and ends in mid-October.

Experimental Animal and Management

A total of 250 goats (180 Red Sokoto and 70 Sahel) of mixed sexes and ages(1 year = 1PPI, 2 years = 2PPI, < 2-3 years = 3PPI and 3.5-4 years above = 4PPI) were used for the study. The animals were randomly selected from Maigatari main livestock market.

Data Collection

Body weight and seven linear body measurements were recorded. They include; body length (BL), heart girth (HG), wither height (WH), testicular length (TL), testicular circumference (TC), udder length (UL) and udder circumference (UC). The measurement was done using a flexible tailors tape, while the wither height was accessed with a measuring stick.

Body Condition Score

Body condition score (BCS) was achieved through palpation of visible bone structure. Animals were palpated over the back bone, loin and rump. Fat condition was smooth to palpate while bone structure with little or no fat cover felt rough to touch. These general assessments were used to place each goat in a category of body condition (very thin = 1, thin = 2, moderate = 3, fat = 4 and obese = 5).

Data Analysis

Data generated were subjected to general linear model (GLM) procedure of SPSS, version 22 (2013). Significantly different means were compared using the least significant differences (LSD). The model utilized was as follow:

$Y_{ijk} = \mathbf{U} + \mathbf{B}_i + \mathbf{S}_j + A_k + \mathbf{e}_{ijk}$

$\mathbf{Y}_{ijk}$ = Observation on dependent variables

U = Common Mean

$\mathbf{B}_i$ = effect of i^{th} breed (1, 2)

$\mathbf{S}_j$ = effect of j^{th} sex (1, 2)

$\mathbf{A}_k$ = effect of k^{th} age (1, 2, 3, 4)

$\mathbf{e}_{ijk}$ = random error term

The relationships among the linear body measurements and body weight were estimated using Pearson's correlation analysis. Stepwise regression procedure was used to find the best linear combination of morphometric traits that can predict body weight. This was performed for male, female and the pooled data. The following linear and multiple regression models were applied:

$Y_1 = a + bx$ - - - - - - (2) simple regression model

$Y_1 = a + b_1x_1 + b_2x_2 + \ldots + b_kx_k$ - - - -(3) multiple regression model

Where Y = dependent variable

a = the Intercept, bs = the slopes and x = independent variable.

RESULTS

Average morphometric traits according to breed, sex and age are presented in Table 1. There was significant effect of breed on BL ($P<0.001$) and WH ($P<0.01$). Sahel goats had lengthier body and higher wither height than Red Sokoto (23.88±0.43 and 24.55±0.41 cm vs 22.45±0.50 and 22.93±0.44 cm). However, non-significant effect of breed on BW, BCS, HG, UL, UC, TL and TC was observed. Significant effect of sex on BW, BL, HG ($P<0.001$) and HG ($P<0.05$) was observed. Females were heavier and had longer body, wider chest and higher wither height than males (25.28±1.00 kg, 24.54±0.41 cm, 26.48±0.40 cm and 24.25±0.38 cm vs 18.51±1.34 kg, 21.80±0.55 cm, 24.47±0.53 cm and 23.24±0.50 cm). Non-significant influence of sex on BSC was however evident. Significant influence of age on BW, BCS, BL, HG, WH ($P<0.001$) and TC ($P<0.01$) was observed. Except TC, goats of 36 - 48 months had the highest for all the significantly affected traits. The correlation coefficients for separate and both sexes are presented in Table 2, 3 and 4. The coefficients observed were in general high, positive and significant except between WH and TL in male and between BW and UC, BCS and BL, WH, UL and UC in females.

Predictive equations of body weight from linear body measurements are shown in Table 5. Of all the body traits recorded, heart girth was observed to have the highest coefficient of determination, while subsequent inclusion of body length in females and pooled data of both sexes yielded better result.

Table 1: Average morphometric traits according to breed, sex and age

		Morphometric traits							
Factor	BW	BSC	BL	HG	WH	UL	UC	TL	TC
Overall mean±S.E.	21.90±1.08	3.14±0.16	23.17±0.42	25.47±0.41	23.74±0.38	3.81±0.48	6.52±1.10	4.86±0.23	7.70±0.23
Breed	NS	NS	**	NS	***	NS	NS	NS	NS
Sahel	22.91±1.11	3.00±0.17	23.88±0.45	25.67±0.44	24.55±0.41	4.14±0.53	7.06±1.19	4.75±0.31	7.52±0.31
Red Sokoto	20.88±1.21	3.20±0.19	22.45±0.50	25.28±0.48	22.93±0.44	3.48±0.60	5.99±1.32	4.97±0.34	7.88±0.33
Sex	***	NS	***	***	*				
Male	18.51±1.34	3.07±0.21	21.80±0.55	24.47±0.53	23.24±0.50	-	-	-	-
Female	25.28±1.00	3.21±0.16	24.54±0.41	26.48±0.40	24.25±0.38	-	-	-	-
Age	***	***	***	***	***	NS	NS	NS	**
Yearlings	17.76±1.25^{b}	2.88±0.20^{c}	20.85±0.51^{b}	23.57±0.49^{b}	22.02±0.46^{b}	3.78±0.71	5.95±1.58	4.44±0.50	7.16±0.50^{b}
12-24 months	19.54±0.90^{b}	2.72±0.14^{c}	21.93±0.37^{b}	24.61±0.36^{b}	23.40±0.33^{b}	3.08±0.58	4.94±1.30	5.95±0.31	6.91±0.31^{b}
24-36 months	27.71±0.86^{a}	3.65±0.14^{b}	25.27±0.35^{a}	27.79±0.34^{a}	25.61±0.32^{a}	4.42±0.44	8.47±0.97	5.53±0.39	9.04±0.38^{a}
36-48 months	27.87±2.67^{a}	3.43±0.42^{a}	25.38±1.10^{a}	28.29±1.10^{a}	25.79±0.98^{a}	5.80±1.23	7.70±2.74	-	-

BW = Bodyweight, BCS = Body condition score, BL= Body length, HG = heart girth, WH= Wither height, TL= Testicular length, TC= Testicular circumference, UL= Udder length and UC= Udder circumference. * = P<0.05, ** = P<0.01, *** = P<0.001, NS = Non-significant and S.E = Standard error

Table 2: Phenotypic correlation coefficients among morphometric traits in male goats

Variable	1	2	3	4	5	6	7
BW (1)	1	0.913**	0.919**	0.983**	0.871**	0.523*	0.791**
BCS (2)		1	0.870**	0.885**	0.830**	0.627**	0.833**
BL (3)			1	0.922**	0.834**	0.445**	0.752**
HG (4)				1	0.872**	0.507**	0.752**
WH (5)					1	0.346^{NS}	0.539*
TL (6)						1	0.774**
TC (7)							1

BW = Bodyweight, BCS = Body condition score, BL= Body length, HG = heart girth, WH= Wither height, TL= Testicular length, TC= Testicular circumference. * = P<0.05, ** = P<0.01 and NS = Non-significant.

Table 3: Phenotypic correlation coefficients among morphometric traits in female goats

Variable	1	2	3	4	5	6	7
BW (1)	1	0.395*	0.818**	0.839**	0.720**	0.371*	0.318NS
BCS (2)		1	0.336NS	0.505**	0.194NS	0.154NS	0.152NS
BL (3)			1	0.754**	0.748**	0.509**	0.436*
HG (4)				1	0.650**	0.487**	0.439*
WH (5)					1	0.419*	0.418*
UL (6)						1	0.847**
UC (7)							1

BW = Bodyweight, BCS = Body condition score, BL= Body length, HG = heart girth, WH= Wither height, UL= Udder length and UC= Udder circumference. * = P<0.05, ** = P<0.01 and NS = Non-significant.

Table 4: Phenotypic correlation coefficients among morphometric traits in both sexes goats

	1	2	3	4	5
BW (1)	1	0.600**	0.898**	0.917**	0.781**
BCS (2)		1	0.598**	0.681**	0.465**
BL (3)			1	0.867**	0.785**
HG (4)				1	0.758**
WH (5)					1

BW = Bodyweight, BCS = Body condition score, BL= Body length, HG = heart girth and WH= Wither height. ** = P<0.01.

Table 5: Prediction equation of body weight from heart girth and body length

Sex	Prediction equation	Coefficient of determination (R^2)
Male	BW = -33.54 + 2.13HG	0.96
Female	BW = -35.04 + 2.28HG	0.69
	BW = -39.50 + 1.40HG + 1.15BL	0.77
Pooled	BW = -39.86 + 2.44HG	0.84
	BW = - 37.81 + 1.50HG + 0.96BL	0.88

BW = Bodyweight, BL= Body length, HG = heart girth. R^2 = Coefficient of determination

DISCUSSION

The facts that breed had effect on BL and WH as observed in the present study agrees with the findings of Guifen *et al*. (2014) who reported a significant variation in morphometric and carcass traits of different genotypes of goats. Similarly, the work of Sanni *et al*. (2018) on different breeds of goats (Kalahari and Sokoto reds, Sahel and WAD) native to Africa also detected a significant variation on morphostructural traits and it has been suggested that the wide variation observed between Sahel and WAD goats could lead to heterotic gain (Zaharaddeen *et al*., 2008). Some authors (Gizaw *et al*., 2007; Agaviezor *et al*., 2012) attributed this variation to inherent genetic potential of each breed, geographical isolation and ecological variation. The significantly higher BSC observed in Red Sokoto goats contradicts the work of Brown and Swan (2012). The results obtained in the present study indicated that morphometric characteristic of Red Sokoto goats was sexually dimorphic in favour of females. This agrees with the work of Ma'aruf *et al*. (2018) in Yankasa sheep breed. Contrary to the present observation, Sowande *et al*. (2008) showed that male sheep were superior to female in live weight and most body measurements. They attributed this differences to genes linked with sex chromosomes. Shuaibu *et al*. (2018 and 2019) also made similar assertion in Yankasa sheep that males had higher values for morphometric traits than females.

The effect of age on live weight and linear body measurements observed in this study agrees with the work of some investigators (Afolayan *et al*., 2006; Mavoule *et al*., 2013) who reported that adult sheep were heavier than the younger ones and had higher values for linear body measurements. Similarly, Mekuriawa *et al*. (2013) reported higher values for morphometric traits in adult sheep compared to the younger and yearling ones and suggested that selective improvement of body weight and conformation traits should be emphasized at this stage (adult). The correlation coefficients observed among the morphometric traits were high, positive and significant. This is in line with the work of Afolayan *et al*. (2006) who reported that live weight was highly correlated with body dimensional traits. The finding of Shuaibu *et al*. (2018) on Yankasa sheep also indicated that the relationship between live weight and linear body measurements were mostly moderate to high which shows that they can predict each other. Working on dairy goat (Saneen), Pesmen and Yerminci (2008) however reported low to moderate relationships among these traits. The highest R^2 observed in this study when BW is predicted from HG conform the report of Tsegaye *et al*. (2013) that this trait (HG) is positively correlated with live weight (LW) and can indicate it with high precision (R^2 = 0.94). Similar report by Tadesse and Gebremariam (2010) in sheep also showed that heart girth predicted LW with higher accuracy of 0.69. Yakubu (2010) attributed this relationship to pleiotropic gene action that exist between the two traits. In the present finding, subsequent inclusion of body length in females and in pooled data of both sexes yielded better values, this agrees with the report of Tadesse and Gebremariam (2010) who observed higher R^2 when BL and WH are included in the predictive equation of BW from HG.

CONCLUSION

I. The results of this study showed that breed had no influence on body weight and most linear body measurements, however, significant effect of sex and age was observed.
II. The correlation coefficients observed among the morphometric traits were in general high, positive and significant.
III. The coefficient of determination observed in the prediction of body weight from height girth was high, but subsequent inclusion of body length in female and the pooled data of sexes yielded better result.
IV. Heart girth predicted body weight with highest precision.

REFERENCES

Ademosun, A. A. (1994). Constraints and prospects for small ruminant research and development in Africa. In: Small Ruminant Research and Development in Africa. *Proceedings of the 2nd Biennial Conference of the African Small Ruminant Research Network, Arcc, Arusha, Tanzania*. 7th – 11th December: 1 - 6.

Agaviezor, B. O., Peters, S. O., Adefenwa, M. A., Yakubu, A., Adebambo, O. A., Ozoje, M. O., Ikeobi, C. O. N., Wheto, M. Ajayi, O. O., Amusan, S. A., Ekundayo, J. O., Sanni, M. T., Okpeku, M., Onasanya, G. O., De Donato, M., Ilori, B. M., Kizikaya, K. and Imumorin I. G. (2012): Morphological and microsatellite DNA diversity of Nigerian indigenous sheep. *Journal of Animal Science and Biotechnology*, **3**: 38 – 44.

Akpa, G. N., Asiribo, O. E., Oni, O. O. and Alawa, J. P. (2001).The influence of non-genetic factors on the shape of lactation curves in Red Sokoto goats. *Nigerian Journal ofAnimal Science*, **72**: 233 - 239.

Afolayan, R. A., Adeyinka, I. A. and Lakpini, C. A. M. (2006). Estimation of live weight from body measurements in Yankasa sheep. *CezchJournal of Animal Science*, **51**(8): 343 – 348.

Brown, D. and Swan, A. (2012). Genetic analysis of adult body weight and condition score in Merino Sheep. *Proceedings, 10th World Congress of Genetics Applied to Livestock Production. Animal Genetics and Breeding Unit, University of New England, Armidale, NSW, Australia*.

Ijomanta, E.O. (2012). Genetic Profile of Morphological Traits and their Association with Production Capacity of Red Sokoto Goat in Katsina State. M.Sc. Thesis, A.B.U, Zaria. Pp: 22 - 26.

Gefu, J. O. (1992). Analysis of factors associated with management of goat in Kano state. *Journal Animal Production Research*, **12** (1): 31 – 39.

Gizaw, S., van Arendonk, J. A. M., Komen, H., Windig, J. J. and Hanotte, O. (2007). Population structure, genetic variation and morphological diversity in indigenous sheep of Ethiopia. *Animal Genetics*, **38**: 621– 628.

Guifen L., Fachun, W., Enliang, S., Xiaomu, L., Xiuwen, T. and Haijian, C. (2014): SNP identification and analysis in part of exon 3 of beef cattle MSTN gene and variation among breeds. *Journal of Animal and Veterinary Advances*, **13**: 174 – 177.

Ma'aruf, B. S., Ibrahim, T., Umar, H. A., Shu'aibu, A. and Okoh, J. J. (2018). Effects of age and sex on body weight, body condition score and biometric traits of Yankasa Sheep. *Nigerian Journal of . Animal Science and Technology***1**(2): 39 – 46.

Mavule, B., Muchenje, V., Bezuidenhout, C. and Kunene, N. (2013). Morphological structure of Zulu sheep based on principal component analysis of body measurements. *Small Ruminant Research*, **111**(1): 23 - 30.

Mekuriawa, S., Mekuriawb, Z., Tayeb, M., Mekuriawb, G., Amanea, A., T. Bimrewa, T. and Haile, A. (2013). Growth performance and linear body measurements of Washera, Farta and their crosses under farmers' management system in Western Highland of Amhara Region. *Scientific Journal of Veterinary Advances*, **2**(9): 132-143.

Otoikhian, C. S. O. and Orheruata, A. M. (2010). The white Bornu goats: Untapped meat-breed in Nigerian small ruminant industry. Klobex Academic Publishers. Bioscience Research Communications 22: 6.

Peacock, C. (2005). Goat: A pathway out of poverty. *Small Ruminant Research*, **60**: 179 - 185.

Pesmen, G. and Yardimci, M. (2008). Estimating the live weight using some body Measurements in Saanen goats. *Archivos de Zootecnica*, **11**: 30 - 40.

Sanni, M. T., Okpeku, M., Onasanya, G. O., Adeleke, M. A., Wheto, M., Adenaike, A. S., Oluwatosin, B. O., Adebambo, O. A. and Ikeobi, C. O. N. (2018). Genetic morphometry in Nigerian and South African Kalahari Red goat breeds. *Agricultura Tropica Et Subtropica*, **51**(2): 51 – 61.

Sownade, O. S. and Sobola, O. S. (2008). Body measurements of West African Dwarf sheep as parameters for estimation of live weight. *Tropical Animal Health and Production*, **40**: 433-439.

Shuaibu, A., Umar, H. A., Ja'afar, A. M., Mujitaba, M. A. and Almajiri, I. (2019). Morphometric characterization of Yankasa sheep in Bauchi South of Bauchi State, Nigeria. *International Journal of Advances in Scientific Research and Engineering,* **5**(2): 34 - 43.

Shuaibu, A., Aliyu, A., Ja'afar, A. M., Ma'aruf, B. S. and Mujitaba, M. A. (2018). Phenotypic characterization of Yankasa sheep in Northern Agricultural Zone of Bauchi State, Nigeria. *Nigerian Journal of Animal Science and Technology*, **1** (2): 36 - 45.

Tadesse, A. and Gebremariam, T. (2010). Application of linear body measurements for live body weight estimation of high sheep in Tigray region, North-Ethipia. *Journal of the Dry land*, **3**(2): 203-207.

Thiruvenkadam, A. K. (2005). Determination of best fitted regression model for estimation of body weight in Kanni Adu kids under farmers' management system. *Livestock Research for Rural Development*, **17** (7): 1-11.

Yakubu, A. (2010). Path coefficients and Path analysis of body weight and biometric traits in Yankasa lambs. *Slovak Journal of Animal Science*,**43**(1): 17-25.

Yakubu, A., Salako, A. E., Imumorin, I. G., Ige, A. O. and Akinyemi, M. O. (2010). Discriminant analysis of morphometric differentiation in the West African Dwarf and Red Sokoto goats. *South African Journal of Animal Science*, **40** (4): 381-387.

Zahraddeen, D., Butswat I. S. R. and Mbap S. T. (2008). Evaluation of some factors influencing growth performance of local goats in Nigeria. *African Journal of Food, Agriculture, Nutrition and Development*, **8**: 464 – 479.